

Beat: Business

WORLD AIDS DAY - FOCUS ON FOOD SAFETY

U.S. FOOD AND DRUG ADMINISTRATION

PARIS - SILVER SPRING, 03.12.2015, 09:12 Time

USPA NEWS - December 1, 2015, is World AIDS Day. The U.S. Food and Drug Administration (FDA) reminds people with HIV/AIDS (and those preparing food for them) about the importance of safe food handling in preventing foodborne illness...

December 1, 2015, is World AIDS Day. The U.S. Food and Drug Administration (FDA) reminds people with HIV/AIDS (and those preparing food for them) about the importance of safe food handling in preventing foodborne illness.

Learn about safe selection and preparation of foods for people with HIV/AIDS in the free booklet Food Safety for People with HIV/AIDS : Download at <http://www.fda.gov/Food/FoodbornellnessContaminants/PeopleAtRisk/ucm312669.htm>

Practicing food safety is critical because the Human Immunodeficiency Virus (HIV) that causes Acquired Immunodeficiency Syndrome (AIDS) can damage, or destroy the body's immune system, making those living with this disease more susceptible to foodborne illness (often called 'food poisoning'). If a person with HIV/AIDS contracts a foodborne illness, he or she is also more likely to have a lengthier illness, undergo hospitalization, or even die. This increased risk underscores the critical role safe food handling plays in managing HIV/AIDS.

Some foods are more risky for people with HIV/AIDS because they are more likely to contain harmful bacteria or viruses. In general, these foods fall into two categories :

- Uncooked fresh fruits and vegetables.
- Certain animal products, such as unpasteurized (raw) milk; soft cheeses made with raw milk; and raw or undercooked eggs, raw meat, raw poultry, raw fish, raw shellfish and their juices; luncheon meats and deli-type salads (without added preservatives) prepared on site in a deli-type establishment.

Anyone who has HIV/AIDS or who prepares food for people with HIV/AIDS should also follow these steps :

- CLEAN: Wash hands and surfaces often. Bacteria can be spread throughout the kitchen and get onto hands, cutting boards, utensils, counter tops, and food.
- SEPARATE: Keep raw meat, poultry, eggs, and seafood and their juices away from ready-to-eat foods.
- COOK to the right temperatures. Use a food thermometer to ensure meat, poultry, seafood, and egg products are cooked to a safe minimum internal temperature to destroy any harmful bacteria. Refer to the chart at: <http://www.fda.gov/downloads/Food/ResourcesForYou/Consumers/UCM462491.pdf>
- CHILL foods promptly. Cold temperatures slow the growth of harmful bacteria. Use an appliance thermometer to be sure the refrigerator temperature is 40 degrees F or below and the freezer temperature is 0 degrees F or below.

Consuming dangerous foodborne bacteria will usually cause illness within 1 to 3 days of eating the contaminated food. However, sickness can also occur within 20 minutes or up to 6 weeks later. Symptoms of foodborne illness include: vomiting, diarrhea, abdominal pain, and flu-like symptoms (such as fever, headache, and body ache).

Source : Food and Drug Administration (FDA)

Ruby BIRD

<http://www.portfolio.uspa24.com/>

Yasmina BEDDOU

<http://www.yasmina-beddou.uspa24.com/>

Article online:

<https://www.uspa24.com/bericht-6635/world-aids-day--focus-on-food-safety.html>

Editorial office and responsibility:

V.i.S.d.P. & Sect. 6 MDStV (German Interstate Media Services Agreement): Ruby BIRD & Yasmina BEDDOU (Journalists/Directors)

Exemption from liability:

The publisher shall assume no liability for the accuracy or completeness of the published report and is merely providing space for the submission of and access to third-party content. Liability for the content of a report lies solely with the author of such report. Ruby BIRD & Yasmina BEDDOU (Journalists/Directors)

Editorial program service of General News Agency:

UPA United Press Agency LTD

483 Green Lanes

UK, London N13NV 4BS

contact (at) unitedpressagency.com

Official Federal Reg. No. 7442619